

Adult and Youth Cigarette Smoking Rates

Adults Youth (grades 9-12)

Sources: Adult 1995 data are BRFSS data reported in *Morbidity and Mortality Weekly Report*, Nov. 8, 1996 (Table 1), www.cdc.gov/mmwr/ preview/mmwrhtml/00044328.htm. Adult 2000 data are BRFSS data reported in *Tobacco Control State Highlights 2002 Report*, www.cdc.gov/ tobacco/statehi/html_2002/state_highlights2002.htm. Youth 1995 (grades 9-12) are YRFSS data reported in *MMWR* Youth Risk Behavior Surveillance–United States, 1995, www.cdc.gov/mmwr/preview/mmwrhtml/00043812.htm. Youth 2000 (grades 9-12) are YRFSS data reported in CDC's *Tobacco Control Program Guidelines and Data State Highlights 2002*, www.cdc.gov/tobacco/statehi/html_2002/ current_youthb.htm. Oregon adults: Oregon Behavioral Risk Factor Surveillance System, 1995 and 2000. Oregon youth: Oregon Public School Drug Use Survey (1996, 2000), Oregon Youth Risk Behavior Survey (1995, 2000), Oregon Healthy Teens Survey (2001).

More Smoking Statistics							
Smoking-attributable deaths, 1999 (per 100,000)	Alaska	Idaho	Montana	Oregon	Washington	Wyoming	USA
	289.6	254.5	294.7	278.6	284.9	318.6	295.5
Smoking-attributable direct medical expenditures, 1998 (per capita	\$215)	\$202	\$246	\$265	\$269	\$221	\$280

Sources: *Investment in Tobacco Control–State Highlights 2001*, www.cdc.gov/tobacco/statehi/pdf_2001/2001statehighlights.pdf. Another source of these data is the CDC database Smoking-attributable mortality, morbidity, and economic costs (SAMMEC): adult SAMMEC and maternal and child health (MCH), www.cdc.gov/tobacco/sammec.